



ANIMAL PREMISES DISINFECTANT CLEANER & DEODORIZER
CONCENTRATE

Reference Sheet

Manufactured by:
Virox Technologies Inc.
Oakville, ON Canada L6H 6R1
Questions? Comments?
1-800-387-7578 | virox.com



FOR VETERINARY USE

For use in animal health facilities and areas. For disinfection and cleaning of hard, non-porous surfaces found in animal premises facilities such as veterinary clinics and offices, animal shelters, kennels, treatment facilities, transportation vehicles, pet grooming facilities, pet boarding facilities, and pet shops.

VIRUCIDAL: 5 MIN. AT 1:40

Avian Influenza A (H3N2) Reassortant Virus
Canine Adenovirus (Strain Toronto A 26/61)
Canine Coronavirus (Strain 1-71)
Canine Distemper Virus
Canine Parainfluenza Virus
Canine Parvovirus
Feline Calicivirus
Feline Infectious Peritonitis Virus
Feline Panleukopenia Virus
Feline Picornavirus (Strain FRV)
Feline Rhinotracheitis Virus (Feline Herpes)
HIV-1 Human Immunodeficiency Virus (HIV) (Strain HTLV-III_B (HIV-1))
Human Coronavirus 229E
Human Parainfluenza Virus 3
Influenza Virus PR8 Strain
Poliovirus Type 1
Pseudorabies Virus
Rhinovirus Serotype 14
Rotavirus (Strain WA)

This product has demonstrated effectiveness against Human Coronavirus (Strain 229E) and is expected to inactivate SARS-CoV-2 (the virus that causes COVID-19).

Proven effectiveness against Poliovirus Type 1, Sabin Strain Type 1 which allows for a Broad-Spectrum Virucide claim against most enveloped and non-enveloped viruses.

This product has demonstrated effectiveness against Poliovirus and is expected to inactivate all Influenza A viruses including 2009 (H1N1) Pandemic Influenza A Virus.

BACTERICIDAL: 5 MIN. AT 1:40

Acinetobacter baumannii
Enterobacter aerogenes
Enterococcus faecalis (Vancomycin Resistant (VRE))
Escherichia coli O157:H7
Klebsiella pneumoniae
Pseudomonas aeruginosa

Salmonella enterica
Staphylococcus aureus
Staphylococcus aureus (Methicillin Resistant (MRSA))

FUNGICIDAL: 5 MIN. AT 1:40

Candida albicans
Trichophyton interdigitale

TUBERCULOCIDAL: 5 MIN. AT 1:40

Mycobacterium terrae

BROAD-SPECTRUM SANITIZING: 30 SEC. AT 1:128

Acinetobacter baumannii
Campylobacter jejuni
Enterococcus faecalis (Vancomycin Resistant (VRE))
Escherichia coli O157:H7
Klebsiella pneumoniae
Pseudomonas aeruginosa
Salmonella enterica
Staphylococcus aureus
Staphylococcus aureus (Methicillin Resistant (MRSA))

DISINFECTANT USE DIRECTIONS

HEAVILY SOILED SURFACES REQUIRE CLEANING PRIOR TO DISINFECTION

USE DILUTIONS MADE EASY

Ratio	Per Litre of Water	Per 4 Litres of Water
1:40	25 mL	100 mL
1:128	8 mL	32 mL

Directions to Mix:

1. Fill empty container with cold water based on the amount required for desired dilution.
2. Carefully add the concentrate into filled container.
3. Gently shake or mix the concentrate and water in the container prior to use.

DO NOT USE UNDILUTED.

Prevail™ Animal Premises Disinfectant Cleaner & Deodorizer Concentrate meets stability testing for up to 30 days after dilution with tap water.

Cleaning, Disinfection and Deodorizing of Equipment and Hard, Non-Porous Surfaces:

1. Pre-clean heavily soiled areas.
2. Dilute 1:40.
3. Apply solution to hard, non-porous surfaces.
4. Allow surface to remain wet for 5 minutes.
5. Let air dry.

Broad-Spectrum Sanitizing on Environmental Surfaces:

1. Dilute 1:128.
2. Apply solution to hard, non-porous surfaces.
3. Allow surface to remain wet for 30 seconds.
4. Wipe dry. No rinsing is required. Food contact surfaces require rinsing with potable water.

Directions for Use in Animal Housing Areas:

1. Remove all animals and their feed from premises, vehicles, and enclosures prior to disinfection.
2. Remove heavy soil such as urine from the surfaces of facilities and fixtures occupied by animals.
3. Empty all feeding and watering appliances.
4. Thoroughly clean all surfaces with detergent or 1:128 dilution of the product and rinse with potable water.
5. Apply 1:40 diluted solution to all hard, non-porous surfaces.
6. Allow surfaces to remain wet for 5 minutes.
7. All surfaces and equipment used in the handling and restraining of animals, and for removing litter, should be immersed in 1:40 diluted solution of the product for 5 minutes. Allow items to dry before re-using.
8. Ventilate closed spaces. Do not house animals or employ equipment until the product has been absorbed, set, or dried.
9. Thoroughly scrub all treated surfaces or objects that come in contact with feed and drinking water such as automatic feeders and waterers with soap or detergent, and rinse with potable water before reuse.

PRECAUTIONARY STATEMENTS

KEEP OUT OF REACH OF CHILDREN

DANGER: Corrosive. Causes severe eye damage. May irritate skin. Harmful if swallowed. Wear goggles or safety glasses, gloves and suitable protective clothing when working with the concentrated product. Wash thoroughly with soap and after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Avoid (i) contact with eyes and skin; (ii) mixing with other cleaning or disinfecting products; and (iii) contamination of food in the application and storage of this product.

FIRST AID

If in contact with eyes, call a physician and flush immediately and thoroughly with water for 15 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. If in contact with skin, flush immediately with water and wash skin thoroughly with soap and water. Obtain medical attention if irritation persists or develops. If ingested, call a physician or a poison control center immediately and do not induce vomiting. TAKE CONTAINER, LABEL OR PRODUCT NAME AND DIN WITH YOU WHEN SEEKING MEDICAL ATTENTION.

TOXICOLOGICAL INFORMATION

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

STORAGE

Avoid storage at elevated temperatures.

DISPOSAL

For information on the disposal of unused, unwanted product and the clean up of spills, contact the Provincial Regulatory Agency or the Manufacturer. Triple rinse the container with water. Dispose of container in accordance with municipal, provincial and federal regulations. Do not reuse container.

DISCLAIMER: You must read and follow the instructions on the label for our product and use the product for the particular applications specified on the product label. To the full extent permitted under applicable law, Virox Technologies Inc. shall not be liable for any damages whatsoever, arising out of or otherwise in connection with any failure to follow those instructions or any use of its products which is not specified on the product's label. The instructions on the product label are not a warranty, express or implied, and Virox Technologies, Inc. expressly disclaims any implied warranties, including any warranty of fitness for a particular purpose. Under applicable law, some of the above waivers or disclaimers of warranty, or exclusions of liability for certain types of damages may not apply to you. In such cases, our liability will be limited to the fullest extent permitted by applicable law.

For relevant patents, visit: virox.com/patents

Accelerated Hydrogen Peroxide® and Design are trademarks of Diversey, Inc.