


### CAGES & KENNELS

#### 4 Steps to Disinfection

**1** Remove animals, food and water bowls and pre-clean by removing any visible soils



**2** Apply the Prevail<sup>™</sup> disinfectant using product of choice<sup>1</sup>



#### CONCENTRATE DILUTION CHART

AREA	DILUTION	CONTACT TIME <sup>1</sup>	DESCRIPTION <sup>1</sup>
Non-Animal Related Areas (Hallways & Offices) testing	1:128 (8 ml / 1 L)	30 seconds	Non-Food Contact Broad Spectrum Sanitization
Daily Disinfection of all Animal Housing Areas and Enclosures	1:40 (25 ml / 1 L)	5 Minutes	Virucide, (including for Canine Parvovirus)
		5 Minutes	Bactericide
High Risk/Quarantine Areas	1:40 (25 ml / 1 L)	5 Minutes	Virucide (including Canine Parvovirus)
Confirmed Disease/Outbreak Conditions	1:40 (25 ml / 1 L)	5 Minutes	Virucide (including Canine Parvovirus), Bactericide, and Fungicide

**3** Ensure surface remains wet for the required contact time<sup>1</sup>



**4** Allow surface to air dry or wipe dry<sup>2</sup>

#### PRODUCT OPTIONS:

##### Prevail<sup>™</sup> Concentrate<sup>1</sup>

- Refer to dilution chart

##### Prevail<sup>™</sup> Ready to Use<sup>1</sup>

- 3 minute contact time

##### Prevail<sup>™</sup> Wipes<sup>1</sup>

- 3 minute contact time

#### SURFACES CHECKLIST:

- Inside of Cage/Kennel
- Front of Cage/Kennel Door (individual bars)
- Cage/Kennel Door Latch

For more information visit:  
**ViroxAnimalHealth.com**

<sup>1</sup> Refer to product label for use instructions and/or reference sheet for complete list of claims and use instructions

<sup>2</sup> Some surfaces may require rinsing